

GEMINI-IFU SPECTROSCOPY OF HH 111 ¹

A. H. CERQUEIRA^{2,a}, M. J. VASCONCELOS^{2,a}, A. C. RAGA^b, J. FEITOSA^{3,a} AND H. PLANA^a

^aLATO-DCET, Universidade Estadual de Santa Cruz

Rod. Jorge Amado, km 16, Ilhéus, BA, Brazil - CEP 45662-900 and

^bInstituto de Ciencias Nucleares, Universidad Nacional Autónoma de México, A.P. 70-543, 04510 D.F., México

Accepted for publication in The Astronomical Journal

ABSTRACT

We present new optical observations of the HH 111 Herbig-Haro jet using the Gemini Multi Object Spectrograph in its Integral Field Unit mode. Eight fields of 5'' x 3.5'' have been positioned along and across the HH 111 jet, covering the spatial region from knot E to L in HH 111 (namely, knots E, F, G, H, J, K and L). We present images and velocity channel maps for the [O I] 6300+6360, H α , [N II] 6548+6583 and [S II] 6716+6730 lines, as well as for the [S II]6716/6730 line ratio. We find that the HH 111 jet has an inner region with lower excitation and higher radial velocity, surrounded by a broader region of higher excitation and lower radial velocity. Also, we find higher electron densities at lower radial velocities. These results imply that the HH 111 jet has a fast, axial region with lower velocity shocks surrounded by a lower velocity sheath with higher velocity shocks.

Subject headings: ISM: Herbig-Haro objects — ISM: jets and outflows — ISM: kinematics and dynamics — ISM: individual (HH 111)

1. INTRODUCTION

HH 111 (discovered by Reipurth 1989) is one of the two most remarkable and better studied Herbig-Haro (HH) jets, the other one being HH 34 (shown to be a jet by Reipurth et al. 1986). The past observational studies of HH 111 include:

- ground based (Reipurth et al. 1992; Podio et al. 2006) and HST (Raga et al. 2002a) high and low resolution long-slit spectra,
- optical (ground based: Reipurth et al. 1992; HST: Hartigan et al. 2001) and IR (Coppin et al. 1998) proper motions,
- evidence of multiplicity of the outflow source (Gredel & Reipurth 1993; Reipurth et al. 1999; Reipurth et al. 2000; Noriega-Crespo et al. 2011),
- discovery of an associated giant jet (Reipurth et al. 1997a),
- detection of an associated, very well collimated molecular outflow (Cernicharo & Reipurth 1996; Nagar et al. 1997; Hatchell et al. 1999; Lefloch et al. 2007)

Numerical simulations of variable jets calculated specifically for modelling the observational properties of

HH 111 were presented by Masciadri et al. (2002) and Raga et al. (2002b).

Optically, the HH 111 system has a one-sided jet that appears at $\sim 15''$ from an obscured source (detected at radio wavelengths by Reipurth et al. 1999), extending to a distance of $\sim 40''$ W from the source. However, a much more symmetric jet/counterjet structure extending down to the position of the source is observed in ground based (Gredel & Reipurth 1994; Davis et al. 1994), HST (Reipurth et al. 1999) and Spitzer (Noriega-Crespo et al. 2011) IR images. At angular scales $> 2'$ from the source, both of the outflow lobes are detected optically, with total extent of $\sim 2^\circ$ for the whole system (see Reipurth et al. 1997a).

The kinematics of the HH 34 jet have been studied spectroscopically (for a limited set of optical emission lines) with full spatial coverage by Beck et al. (2007) and Rodríguez-González et al. (2012). Our present paper describes similar observations (i.e., high resolution spectroscopy with full spatial coverage), but for the HH 111 jet. In our Integral Field Unit (IFU) Gemini South spectra the [S II] 6730, 6716; H α ; [N II] 6583, 6548 and [O I] 6360, 6300 are detected, and have a high enough signal-to-noise so that velocity channel maps can be generated.

The paper is organized as follows. In section 2 we describe the observations and the data reduction. The results are described in section 3. Finally, the work is summarized in section 4.

2. OBSERVATIONS AND DATA REDUCTION

2.1. The observations

HH 111 was observed at GEMINI North observatory on Oct 2007 and Nov 2010, using the GMOS instrument in IFU mode (Allington-Smith et al. 2002) under the program GN-2007B-Q-9. We used the IFU in the single slit mode with the R831.G5302 grating, giving a $R = 4396$ resolution at 7570Å. In the single slit mode, the IFU field of view is 5'' x 3''5, each lens covering 0''2 on the sky.

hoth@uesc.br

¹ Based on observations obtained at the Gemini Observatory, which is operated by the Association of Universities for Research in Astronomy, Inc., under a cooperative agreement with the NSF on behalf of the Gemini partnership: the National Science Foundation (United States), the Science and Technology Facilities Council (United Kingdom), the National Research Council (Canada), CONICYT (Chile), the Australian Research Council (Australia), Ministério da Ciência, Tecnologia e Inovação (Brazil) and Ministerio de Ciencia, Tecnología e Innovación Productiva (Argentina).

² On a sabbatical leave at IPAG/UJF, Grenoble, France.

³ INCT-A scholarship

TABLE 1
TARGET COORDINATES AND OFFSETS

Target	RA (h:m:s)	DEC (°:′:″)	Δp (″)	Δq^a (″)
IRAS 05491+0247 ^b	05:51:46.25	02:48:29.5	-	-
field 1	05:51:43.7	02:48:34.1	-	-
field 2			0	4.4
field 3			0	8.8
field 4			2.9	5.6
field 5			-2.9	5.6
field 6			0	-4.4
field 7			0.214	-9.07
field 8			0.214	-13.43

^aThe offsets Δp and Δq are given in arcsec, and refers to displacements perpendicular (p) and parallel (q) to the HH 111 PA; here defined as $\sim 83^\circ$. Field 1 is the reference to the displacement shifts.

^bCoordinates of the HH 111 IRS (Reipurth et al. 2000).




FIG. 1.— Sketch of the eight fields observed with the GMOS-IFU superimposed on the $H\alpha$ pre-image of HH 111 taken with the Gemini North Telescope. The observed fields are labeled by numbers from 1 (centered on the brilliant knot J) to 8 (finishing the mosaic in the double knot E). The knots inside the HH 111 jet are labeled by capital letters from E to L, following the nomenclature provided by Reipurth (1989) and Raga et al. (2002a). The offsets between each field are indicated in Table 1. The N-E axes are indicated in the Figure, as well as a distance scale.

The observations has made under exceptional seeing conditions. Using the R-band images, the seeing estimates range from $0''.44$ to $0''.51$ for the first epoch observations (2007) and from $0''.55$ to $0''.60$ for the second one (2010).

$H\alpha$ narrow-band filter pre-images (on and off) were also taken (on August 22nd, 2007) to position the IFU field with higher accuracy. In order to cover the target, we observed the 8 fields shown in Figure 1. Each




FIG. 2.— *Left*: $H\alpha$ sub-cube of field 8 without any filtering process ($3''.5 \times 5''.0$). *Middle*: The sum of the first two eigenvectors, after PCA analysis of the data cube in the spectral region defined by $H\alpha$, but excluding the emission. *Right*: First image (left panel) minus the low-frequency noise (middle panel). The results show that the instrumental fingerprint has been removed from the original data, without any introduction of artificial structures.

field was observed with a total exposure of 1200s (three 400s exposures, which were then median averaged to reduce the cosmic ray contamination). Fields 1 to 6 were observed on October 19th 2007 and fields 7 and 8 on November 16th 2007. Fields 1 to 6 were re-observed on November 2nd 2010, due to bad CCD settings during the original observations. Overlaps between the different fields have been set to $0''.4$ (corresponding to two rows of IFU lenses).

Table 1 gives the coordinates of the HH 111 IRS outflow source (IRAS 05491+0247; see, Rodríguez & Reipurth 1994; Rodríguez et al. 1998; Reipurth et al. 2000) and the exact offsets of the GMOS-IFU observed fields. The offsets are in arcsec and are taken with respect to the center of field 1 (center of knot J, see Figure 1). The offsets are perpendicular (Δp) and parallel (Δq) to the HH 111 outflow axis, which is at a PA= -83.94° position angle (Raga et al. 2002a).

2.2. Data reduction

Data have been reduced using special reduction package provided by Gemini Staff⁴ using the IRAF reduction package⁵. All raw images have been bias corrected, trimmed and flat fielded. Flatfield images have also been used to located the positions of the 1500 lenses on the frame. Twilight images were used to estimate the grating response and after the extraction of spectra for each lens, each one has been corrected using this normalized response. Arcs, from the CuAr lamp, have also been taken for the wavelength calibration. Using the bright OI night sky line at $5577.338 \text{ \AA}$, we have estimated the wavelength calibration accuracy to $0.1 \text{ \AA}$. The last step of the reduction was the sky subtraction using the field located at $1'$ from the science field. Finally the data cube has been created with the GFCUBE task using a spatial resampling of $0.1''$ per pixel. The total spectral coverage goes from $4835.896 \text{ \AA}$ to $6957.800 \text{ \AA}$. The spectral sampling is $0.339 \text{ \AA}$ per spectral pixel. We then extracted sub-cubes near the $H\alpha$, $[\text{O I}]\lambda\lambda 6300,6364$, $[\text{N II}]\lambda\lambda 6548,6584$ and $[\text{S II}]\lambda\lambda 6716,6731$ emission lines, for each the eight observed fields.

Further image treatment has been conducted using the

⁴ <http://www.gemini.edu/node/10795>

⁵ Image Reduction and Analysis Facility is a software developed by the National Optical Astronomy Observatory - iraf.noao.edu




FIG. 3.— *From left to right:* Mosaics of [S II] $\lambda 6716 + \lambda 6731$ for the integrated line profile (left), and for three different radial velocities intervals: high velocity (HVC, medium velocity (MVC) and low velocity (LVC) components. The intervals are defined by: $v_{rad} < -100$ km s $^{-1}$ for HVC, $-100 < v_{rad} < -70$ (in km s $^{-1}$) for the MVC and $v_{rad} > -70$ km s $^{-1}$ for the LVC. The coordinate axes are in arcsecs (the origin coincides with the position of the driving source) and the colorbar are in arbitrary units.




FIG. 4.— The same as in Figure 3 for H α .

pipeline developed by Menezes et al. (2014; see also Ricci, Steiner & Menezes 2011). It consist in i) a spatial filtering process to remove high-spatial frequency noise, ii) a Principal Component Analysis (PCA) to remove instrumental spatial fingerprints and iii) a Richardson-Lucy deconvolution process. In order to remove the high spatial frequency noise, we first calculate discrete Fourier transforms of the images resulting from each data cube, and apply a low-pass band filter in the frequency space

(Gonzalez & Woods 2002). To this effect we follow the procedure described in Menezes, Steiner & Ricci (2014). In particular, we use a Butterworth $H(u, v)$ filter of order $n = 6$ (see equation 7 in Menezes et al. 2014). An inverse Fourier transform is then applied to the filtered data. We should mention that similar noise filtering process has been already applied in the context of long slit spectroscopy of HH jets (Raga & Mateo 1998). The high-frequency cleaned data still show low-frequency




FIG. 5.— The same as in Figure 3 but for $[\text{N II}]\lambda 6548 + \lambda 6583$.




FIG. 6.— The same as in Figure 3 but for $[\text{O I}]\lambda 6300 + \lambda 6360$.

noise which we remove using the Principal Component Analysis (PCA; see Ricci, Steiner & Menezes 2011 and Steiner et al. 2009 for a detailed discussion and method presentation). The scope of this paper is not to present this technique. However, it is worth to mention briefly how it works. The PCA technique allows us to describe the data cube as a linear combination of an orthogonal basis, which are the eigenvectors of a 2D covariance matrix. Such a covariance matrix is obtained after scaling all the intensities in the 3D data cube with the transfor-

mation:

$$\beta = \mu(i - 1) + j \quad (1)$$

in a data cube with the spatial dimensions $n = \mu \times \nu$: $1 \leq i \leq \mu$, $1 \leq j \leq \nu$. For each spaxel intensity, $I_{ij\lambda}$, one defines a 2D intensity matrix $\mathbf{I}_{\beta\lambda}$, which will be the subject of the PCA analysis. Its covariance matrix is calculated, and the eigenvectors with their associated variances are obtained (e.g., Steiner et al. 2009). The higher its variance, the more representative of the




FIG. 7.— Low-velocity and high-velocity components mosaic subtraction in $H\alpha$ emission. *Left*: $LVC - 2 \times HVC$. *Right*: $LVC - 4 \times HVC$.

data is a given eigenvector. As the orthogonality is ensured, we can compose a tomogram, which is an image that represents the projection of the original data onto a selected eigenvector (or eigenvectors). In Figure 2 we show a specific observed field (field 8 in Figure 1), for a given emission line ($H\alpha$), before any filtering process (a), after the filtering process but extracting the emission line from data (b), and the (a)-(b) subtraction in (c). We should mention that the tomogram in (b) was obtained by adding the first two eigenvectors (without the emission line), which we assume to correspond to the low-frequency noise, or the instrumental fingerprint, as described in Ricci et al. (2011).

Finally, we have applied a Richardson-Lucy deconvolution procedure with a Gaussian PSF, with a constant FWHM of $0''.44$ for fields 7 and 8, and a constant FWHM of $0''.55$ for the fields 1 to 6, as suggested by the seeing in our observations. Six interactions has been used in order to deconvolve the original data with the PSF (see equation 10 in Menezes et al. 2014). The whole procedure (spatial re-sampling, low-noise filtering process, instrumental fingerprints removal technique and Richardson-Lucy deconvolution) is explained in detail in Menezes et al. (2014), who has developed the technique⁶.

3. RESULTS

3.1. Images and velocity channel maps

We present the spatially resolved line profiles as “velocity channel maps” in three velocity intervals:

- HVC: a high (negative) velocity channel, with radial velocities⁷ $v_{rad} < -100 \text{ km s}^{-1}$,
- MVC: a medium velocity channel, with $-100 \text{ km s}^{-1} < v_{rad} < -70 \text{ km s}^{-1}$,
- LVC: a low velocity channel, with $v_{rad} > -70 \text{ km s}^{-1}$.

⁶ The routines are available electronically at the following address: [http://www.astro.iag.usp.br/~sim\\$PCAtomography/](http://www.astro.iag.usp.br/~sim$PCAtomography/).

⁷ The velocities has not been corrected for the LSR velocity of $+23 \text{ km s}^{-1}$ Reipurth 1989.

We also present an image (for each spectral line) which consists of an addition of these three velocity channel maps. The channel maps and the images have been computed for mosaics composed of the 8 observed IFU fields (see Figure 1), and cover the emission of the HH 111 jet at distances $\approx 24'' \rightarrow 50''$ from the outflow source. The mosaics have been built taking into account the offsets (given in Table 1).

In Figures 3, 4, 5 and 6 we show the images and velocity channel maps for $[S \text{ II}]\lambda 6716 + \lambda 6731$, $H\alpha$, $[N \text{ II}]\lambda 6548 + \lambda 6584$ and $[O \text{ I}]\lambda 6300 + \lambda 6360$, respectively. In the $[S \text{ II}]$ image (left frame of figure 3) we see knots E through L, which can be clearly recognized comparing the image with the identifications of Reipurth et al. (1992) and Reipurth et al. (1997b).

In the channel maps of all of the observed lines, it is clear that there is a general trend of increasing jet width as a function of decreasing velocity (i.e., in all of the observed emission lines the jet becomes narrower as we go from the HVC to the MVC and to the LVC, see Figures 3-6). This result is in qualitative agreement with the observations of Riera et al. (2001), who found velocities decreasing away from the outflow axis in two long-slit spectra cutting the HH 111 jet at the approximate positions of knots D (not detected in the present observations) and F.

3.2. Width vs. radial velocity dependence

In order to illustrate the broadening of the jet at lower (i.e., less negative) radial velocities, in Figure 7 we show the LVC–HVC channel map subtraction for the $H\alpha$ line. This subtraction map shows that at all positions along the observed region of the HH 111, the high radial velocity emission is restricted to a narrow, central region of the jet cross section.

One can quantify the width vs. radial velocity dependence by measuring the FWHM of the emission on the HVC, MVC and LVC channel maps. For all of the emission lines, we measure the widths on the three channel maps at the positions of knots E, F, G, H, J and L, integrating the emission in bins of $1''$ along the jet bin (centered at the positions of the knots). The results of this procedure are shown in Figure 8, which shows the following features⁸

- there is a systematic increase of width as a function of decreasing radial velocity for all the knots, and in all emission lines,
- with the exception of the very broad knot E, there is a general trend of increasing width vs. distance from the source,
- in the lower excitation lines ($[O \text{ I}]$ and $[S \text{ II}]$), the widths are somewhat lower than in the higher excitation lines ($H\alpha$ and $[N \text{ II}]$).

This last effect has been noted by Reipurth et al. (1997), who note that the $H\alpha$ emission is broader than

⁸ The determination of the FWHM for the knots G1, J, K1 and L, using the $[N \text{ II}]$ and $[O \text{ I}]$ lines show high uncertainties due to their low S/N (see Figures 5 and 6). In a few cases (knots G1, H and K1; see Figure 8) we were not able to determine the $[N \text{ II}]$ FWHM for some (if not all) velocity components.




FIG. 8.— FWHM (in arcsecs) of the emission on the HVC, MVC and LVC channel maps for all of the emission lines (indicated on the top of the figure), at the positions of knots E, F, G, H, J and L (indicated on the right of the figure). The FWHM has been obtained integrating the emission in bins of $1''$ along the jet (centered at the positions of the knots) and fitting a Gaussian to each knot longitudinal profile. For [N II] line we found it difficult to adjust a profile due to low S/N in some knots and for some velocity components (see also Figure 5). For these cases, the FWHM appears in the histogram as an zero. The H α and [N II] histograms for the knot L has the FWHM axis divided by a factor of 2.

the [S II] emission (see their Figure 6, in which they present an H α -[S II] subtraction map). In our observations, we see that in general the [N II] emission has widths comparable to the H α widths, and that the [O I] emission has narrower widths, comparable to the ones of the [S II] emission.

3.3. The [S II] 6716/6731 line ratios

In Figure 9 we present the spatial distribution of the [S II] λ 6716/[S II] λ 6731 line ratio for the integrated line profiles and for the HVC, MVC and LVC maps. This ratio gives the electron density (n_e) of the medium Osterbrock (1989). For calculating n_e we assume a temperature of 10^4 K, consistent with the temperature of $\sim 1.3 \times 10^4$ K found by Podio et al. (2006) for the HH 111 jet.

Superimposed on the spatial distribution of the electron density are the (logarithmic spaced) iso-contours of the [S II] λ 6716 + λ 6731 integrated intensity. The integrated map (left) shows that the electron density ranges from $\sim 1.5 \times 10^3 \text{ cm}^{-3}$ at knot E (at ~ 26 arcsec from the

outflow source) up to $5 \times 10^3 \text{ cm}^{-3}$ in knot F (at $\sim 2''$ from knot E; see Figure 9). It then drops to $\sim 2 \times 10^3 \text{ cm}^{-3}$ between knots F and H. In knot H, the electron density has a strong peak of $n_e \sim 3.5 \times 10^3 \text{ cm}^{-3}$.

Interestingly, the electron density maps of the three velocity channels show systematically increasing densities for decreasing radial velocities. This can be clearly seen at the positions of knots E, F and J. Also, an increase in n_e can be seen in the region between knots F and G when going from the MVC to the LVC electron density maps (see the two frames on the right of Figure 9).

3.4. Knot E

We now focus on the interesting structure of knot E (at $\sim 26''$ from the source, see Figures 3-7). The structures observed in the images and channel maps of this knot are shown in Figure 10.

We see that in the H α and [N II] images (2nd and 3rd rows of the far left column of Figure 10) knot E shows




FIG. 9.— From left to right: Mosaics of $[\text{S II}]\lambda 6716/[\text{S II}]\lambda 6731$ line ratio for the integrated line profile, and for three different radial velocities intervals: high velocity (HVC), medium velocity (MVC) and low velocity (LVC) components. The intervals are defined by: $v_{\text{rad}} < -100 \text{ km s}^{-1}$ for HVC, $-100 < v_{\text{rad}} < -70$ (in km s^{-1}) for the MVC and $v_{\text{rad}} > -70 \text{ km s}^{-1}$ for the LVC. The coordinate axes are in arcsecs and the colorbar indicates the electron density in cm^{-3} .

two side-by-side peaks. These peaks have been observed in previous ground based (Reipurth et al. 1992) and HST (Reipurth et al. 1997) $\text{H}\alpha$ images of HH 111. Most interestingly, the two side-by-side peaks are well separated in the LVC $\text{H}\alpha$ and $[\text{N II}]$ maps (right column of Figure 10), approach each other in the MVC map, and merge into a single, central peak in the HVC map. Therefore, we find that the two side-by-side peaks (previously observed in $\text{H}\alpha$ images, see Reipurth et al. 1997) actually enclose a fainter, higher radial velocity emission region.

In the lower excitation $[\text{S II}]$ and $[\text{O I}]$ lines, knot E shows two low intensity side-by-side peaks only in the LVC maps (1st and 4th rows of the far right column of Figure 10), and these peaks are not present in the MVC and HVC maps. The images of knot E in these lines (1st and 4th rows of the far left column of Figure 10) show a conical structure which resembles the MVC emission.

4. SUMMARY

We have obtained IFU spectra of the HH 111 jet at distances of $24 \rightarrow 49''$ from the outflow source. From the spectra, we obtain position-velocity cubes for the $[\text{S II}] 6716, 6731$; $\text{H}\alpha$; $[\text{N II}] 6548, 6583$ and $[\text{O I}] 6300, 6360$ emission lines.

We find a number of interesting results:

- we find that for all emission lines we have an increase in the observed width of the jet for decreasing radial velocities. This effect can be seen directly in the velocity channel maps (Figures 3-6), in subtractions of pairs of velocity channel maps (Figure 7) or more quantitatively in the widths measured for the jet knots (Figure 8). A broadening in the HH 111 jet for decreasing velocities has been previously observed by Riera et al. (2001), and a qualitatively similar effect is seen in the CO emission (but at widths in excess of $\sim 10''$, see Lefloch et al. 2007),

- we find that the HH 111 jet is narrower in the lower ($[\text{S II}]$ and $[\text{O I}]$) than in the higher ($\text{H}\alpha$ and $[\text{N II}]$) excitation lines at the vicinity of some knots (namely, knots E, F K1 and L). This effect has been previously seen in the $\text{H}\alpha$ and $[\text{S II}]$ HST images of Reipurth et al. (1997). These previous observations suggested that we might be seeing a sheath of non-radiative shocks (seen only in $\text{H}\alpha$) surrounding the HH 111 jet beam, but the detection of broad $[\text{N II}]$ emission indicates that the shocks in the sheath are indeed radiative,
- from the $[\text{S II}] 6716/6731$ line ratio we systematically find larger electron densities at lower radial velocities (see Figure 9),
- the “twin peak” structure of knot E is clearly seen in the $\text{H}\alpha$ and $[\text{N II}]$ images, but these two peaks merge into a central peak in the $[\text{O I}]$ and $[\text{S II}]$ images. In all of the lines, the twin peaks are seen at low radial velocities (see the LVC maps of Figure 10), but they approach and merge into a single peak at higher radial velocities.

These characteristics can be qualitatively explained in terms of an “internal working surface” model, in which the knots are produced as a result of an outflow velocity time-variability. In such a model, the variability produces pairs of shocks that travel down the jet beam, producing a high pressure region that ejects material into the cocoon of the jet. The observations of larger widths for the higher excitation emission would then be interpreted as higher velocity shocks produced by the material in this sideways ejection against a slower moving cocoon. The low excitation emission would be interpreted as coming from lower velocity shocks within the jet beam (produced by the outflow velocity variability).

The shock structure of internal working surfaces was described by Raga et al. (1990), and numerical simula-




FIG. 10.— *From left to right:* Integrated emission, HVC, MVC and LVC for (from top to bottom) [S II], H α , [N II] and [O I] for field 8 which contains the knot E. The axes are in arcsec and indicates the distance from the driving source. The maps are in arbitrary units (as seen at the colorbars).

tions have been done for reproducing $H\alpha$ images (Raga et al. 2002b) and long-slit spectra (Masciadri et al. 2002) of the HH 111 jet. The data presented here clearly provide a substantial increase in the observational constraints, which should be a challenging test for future models of jet knots produced by an outflow variability or by other mechanisms (see e.g., Micono et al. 1998). In particular, our data will give interesting constraints on the nature of the cross section of the outflow (Raga et al. 2011) or of the possible presence of two distinct components (Teşileanu et al. 2014).

We can speculate that MHD effects might be present, and might help to shape the morphology of the HH 111 emission knots as well as their line profiles. It is worth to note that recent and sophisticated MHD models predict that beam recollimation (by the hoop stress) plays an important role in order to enhance the brightness near the jet axis, as well as to accelerate the inner jet beam, making it faster than the surrounding jet material. This general result was obtained by Teşileanu et al (2014), Hansen, Frank & Hartigan (2014) and Staff et al. (2015), in spite of the differences in the initial setup in these works. In Hansen, Frank & Hartigan (2014), a pure toroidal magnetic field is used to study the jet propagation, while the disk wind solution is incorporated in the models from Staff et al (2014) and Teşileanu et al. (2014), who actually introduced a two component (stellar jet plus disk wind) flow.

Another feature shown by some HH jets (including some of the knots studied here) is an increase of the FWHM of the jet beam with distance from the driving

source. This feature has been predicted in the models presented by Teşileanu et al. (2014) and compared with available data from RW Aur, HH 30 and HL Tau, while the solutions presented in Staff et al. (2014) were compared with data from DG Tau, HN Tau, RW Aur and UZ Tau. In both cases, the distances covered by the simulations were suitable for the study of the microjets (~ 100 AU). Whether or not these effects are still important farther away from the driving source (the knot E is at ~ 10000 AU from the VLA 1 source) is an issue that should be addressed properly in future work.

We would like to thank the referee for his/her suggestion. We are grateful to R. Carrasco and B. Miller (from Gemini South telescope), and R. Schiavon (Liverpool John Moores University) for help us with data reduction and scripts. We are thankful to T. Ricci and J. Steiner for enlightening discussions about the PCA technique and its applications to data-cube, and for providing the pipeline for data reduction. AHC, MJV and HP thanks Cnpq/CAPEs for financial support using the PROCAD project (552236/2011-0) and CAPEs/CNPq Science without Borders program, under grants 2168/13-8 (AHC) and 2565/13-7 (MJV). J. Feitosa was fully supported by a INCT-A scholarship. AR acknowledges support from CONACyT grants 101356, 101975 and 167611 and the DGAPA-UNAM grants IN105312 and IG100214. This paper has been submitted during our (AHC and MJV) sabbatical leave in Grenoble (IPAG/UJF), and we are very thankful to J. Bouvier and J. Ferreira for their warm hospitality.

Facilities: Gemini (GMOS).

REFERENCES

- Allington-Smith, J., Graham, M., Content, R., Dodsworth, G., Davies, R., Miller, B. W., Jørgensen, I., Hook, I., Crampton, D., Murowinski, R. 2002 PASP, 114, 892
- Beck, T. L., Riera, A., Raga, A. C., & Reipurth, B. 2007, AJ, 133, 1221
- Cernicharo, J., & Reipurth, B. 1996, ApJ, 460, L57
- Coppin, K. E. K., Davis, C. J., Micono, M. 1998, MNRAS, 301, L10
- Davis, C. J., Mundt, R., & Eislöffel, J. 1994, ApJ, 437, L55
- Gonzalez, R.C., and Woods, R.E. 2002, in *Digital Image Processing*, 2nd Edition (Upper Saddle River, NJ: Prentice Hall)
- Gredel, R., & Reipurth, B. 1993, ApJ, 407, L29
- Gredel, R., & Reipurth, B. 1994, A&A, 289, L19
- Hansen, E.C., Frank, A., & Hartigan, P. 2014 (arXiv:1412.6489)
- Hatchell, J., Fuller, G. A., & Ladd, E. F. 1999, A&A, 346, 278
- Hartigan, P., Morse, J., Reipurth, B., Heathcote, S., & Bally, J. 2001, ApJ, 559, L157
- Lefloch, B., Cernicharo, J., Reipurth, B., Pardo, J. R., & Neri, R. 2007, ApJ, 658, 498
- Masciadri, E., Velázquez, P. F., Raga, A. C., Cantó, J., Noriega-Crespo, A. 2002, ApJ, 573, 260
- Menezes, R.B., Steiner, J.E. & Ricci, T.V. 2014, MNRAS, 438, 2597
- Micono, M., Massaglia, S., Bodo, G., Rossi, P., & Ferrari, A. 1998, A&A, 333, 1001
- Nagar, N. M., Vogel, S. N., Stone, J. M., Ostriker, E. C. 1997, ApJ, 482, L195
- Noriega-Crespo, A., Garnavich, P. M., & Raga, A. C. 1993, AJ, 106, 1133
- Noriega-Crespo, A., Raga, A. C., Lora, V., Stapelfeldt, K. R., & Carey, S. J. 2011, ApJ, 732, L16
- Osterbrock, D.E. 1989, *Astrophysics of Gaseous Nebulae and Active Galactic Nuclei* (Mill Valley: Univ. Science)
- Podio, L., Bacciotti, F., Nisini, B., Eislöffel, J., Massi, F., Giannini, T. and Ray, T.P. 2006, A&A, 458, 204
- Raga, A. C., Cantó, J., Binette, L., & Calvet, N. 1990, ApJ, 364, 601
- Raga, A.C., and Mateo, M. 1988, RMxAA, 16, 13
- Raga, A.C., Noriega-Crespo, A., Kajdic, P., De Colle, F., López-Cámara, D., Esquivel, A. 2011, RMxAA, 47, 277
- Raga, A.C., Noriega-Crespo, A., Reipurth, B., Garnavich, P.M., Heathcote, S., Böhm, K.H. and Curiel, S., 2002a, ApJ, 565, L33
- Raga, A.C., Velázquez, P. F., Cantó, J., Masciadri, E. 2002b, A&A, 395, 647
- Reipurth, B. 1989, Nature, 340, 42
- Reipurth, B., Bally, J., and Devine, D. 1997, AJ, 114, 2708
- Reipurth, B., Bally, J., Graham, J. A., Lane, A. P., Zealy, W. J. 1986, A&A, 164, 51
- Reipurth, B., Bally, J., Devine, D. 1997a, AJ, 114, 2708
- Reipurth, B., Hartigan, P., Heathcote, S., Morse, J. A., & Bally, J. 1997b, AJ, 114, 757
- Reipurth, B., Raga, A. C., Heathcote, S. 1992, ApJ, 392, 145
- Reipurth, B., Yu, K.C., Heathcote, S., Bally, J., and Rodríguez, L.F. 1999, A&A, 352, L83
- Reipurth, B., Yu, K.C., Heathcote, S., Bally, J., and Rodríguez, L.F. 2000, AJ, 120, 1449
- Reipurth, B., Raga, A.C., and Heathcote, S. 1992, ApJ, 392, 145
- Ricci, T., Steiner, J., and Menezes, R.B. 2011, ApJ, 734, L10
- Riera, A., López, R., Raga, A. C., Anglada, G., & Estalella, R. 2001, RMxAA, 37, 147
- Rodríguez, L.F. and Reipurth, B. 1994, A&A, 281, 882
- Rodríguez, L.F., Reipurth, B., Raga, A.C. and Cantó, J. 1994, RMxAA, 34, 69
- Rodríguez-González, A., Esquivel, A., Raga, A. C., Cantó, J., Riera, A., Curiel, S., & Beck, T. L. 2012, AJ, 143, 60
- Staff, J.E., Koning, N., Ouyed, R., Thompson, A., Pudritz, R.E. 2015 MNRAS, 446, 3975

Steiner, J., Menezes, R.B., Ricci, T., and Oliveira, A. 2009,
MNRAS, 396, 788
Teşileanu, O., et al. 2014, A&A, 562, A117

Vasconcelos, M.J., Cerqueira, A.H., Plana, H., Raga, A.C. and
Morisset, C. 2005, AJ, 130, 1707